



**PERMIT TO ALLOW MINOR USE
OF A REGISTERED AGVET CHEMICAL PRODUCT**

**FOR CONTROL OF ANTHRACNOSE AND PANICLE SHOOT BLIGHT IN
PISTACHIOS**

PERMIT NUMBER – PER80760

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 10 SEPTEMBER 2015 to 31 AUGUST 2020

Permit Holder:

PISTACHIO GROWERS AUST INC
c/- AgAware Consulting Pty Ltd
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Product to be used:

BRAVO FUNGICIDE

Plus ALL OTHER REGISTERED PRODUCTS

Containing: 720 g/kg CHLOROTHALONIL as the only active constituent.

Directions for Use:

Crop	Diseases	Rate
PISTACHIOS	Anthracnose (<i>Colletotrichum</i> spp.)	160 – 210 mL product / 100 L
	Panicle and shoot blight (<i>Botryosphaeria</i> spp.)	OR 1.6 – 2.1 L product / ha

Critical Use Comments:

- Chlorothalonil based products should be applied as a protective treatment prior to the onset of disease development.
- Make first foliar spray application during early bloom. Repeat application if conditions remain favourable for disease development.
- Apply using orchard airblast/mister sprayer in sufficient water to obtain uniform coverage.
- DO NOT apply more than five (5) applications per crop, with a minimum re-treatment interval of 10 days between consecutive applications.
- Use the higher application rate when conditions favour disease development.
- Use disease prediction models (i.e. Leaf Wetness Model developed in California) to determine favourable disease conditions.
- DO NOT mix chlorothalonil based products with spraying oil, wetting agent or other EC fungicide formulations, as crop damage may occur.
- Alternate with fungicide spray treatments from other chemical (Mode of Action) groups. Multiple fungicides are required from different chemical groups to manage the resistance risk.

Withholding Period:

DO NOT harvest for 14 DAYS after application.

Jurisdiction:

ALL STATES, except VIC.

Note: Victoria is not included in this permit, as their *Control-of-Use* legislation means a permit is not required to legalise this off-label use in that state.

Additional Conditions:

This PERMIT provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

PERSONS who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the DETAILS and CONDITIONS of this permit.

Export of produce:

To allow treated produce to be supplied or otherwise made available for human consumption a Temporary Maximum Residue Limit (TMRL) has been established for Chlorothalonil covering the commodity Pistachios at T0.1 mg/kg. This TMRL applies only to produce marketed and consumed in Australia. If treated produce is to be exported, due account should be taken of the residue definition and residue limits/import tolerances of importing countries and that any residues must not exceed those requirements of the importing country.

Issued by the Australian Pesticides and Veterinary Medicines Authority